

DETERMINATION OF THE LOW Q^2 EVOLUTION OF THE BJORKEN INTEGRAL

A. DEUR

*Thomas Jefferson National Accelerator Facility
12000 Jefferson Avenue, Newport News, VA23606, USA
E-mail: deurpam@jlab.gov*

We report on an experimental determination of the Q^2 -dependence of the Bjorken sum using data from Jefferson Lab Hall A and Hall B in the range $0.16 < Q^2 < 1.1$ GeV². A twist analysis is performed. Overall, the higher twist corrections are found to be small due to a cancellation between the twist 4 and 6 terms.

1. The GDH and Bjorken Sum Rules

A main reason to study the generalized GDH sum is to understand the transition from the hadronic to the partonic descriptions of the nucleon. The generalized GDH sum is in principle calculable at any Q^2 , which makes it an ideal tool to study the hadronic to partonic transition. This topic was covered in the symposium¹. However, the validity domains of the available chiral perturbation theory (χ PT) and pQCD calculations do not overlap. Lattice QCD should provide the bridge between the two domains but no calculation is available yet.

The Bjorken sum rule² has been a cornerstone of polarized pQCD studies. At leading twist, it reads:

$$\int_0^{1^-} (g_1^p - g_1^n) dx = \frac{g_a}{6} \left[1 - \frac{\alpha_s}{\pi} - 3.58 \left(\frac{\alpha_s}{\pi} \right)^2 - 20.21 \left(\frac{\alpha_s}{\pi} \right)^3 + \dots \right] \quad (1)$$

where g_a is the nucleon axial charge. The connection between a generalized GDH sum and the Bjorken sum was made by M. Anselmino *et al.*³. More recently, X. Ji and J. Osborne made the reason for the connection clear: the GDH and the Bjorken sum rules are two particular cases of a more general sum rule⁴. The extended GDH sum rule, as generalized in Ref.⁴, links the first moment of the spin structure functions to the spin dependent forward Compton scattering amplitudes. Hence, the relation between the

generalized GDH and Bjorken sums is:

$$\int_0^{1-} g_1^p - g_1^n dx = \frac{Q^2}{16\pi^2\alpha} (\text{GDH}^p(Q^2) - \text{GDH}^n(Q^2)) \quad (2)$$

Considering this $p - n$ flavor non-singlet quantity yields many advantages.

- (1) At large Q^2 , we have a sum rule (the Bjorken sum rule) without hypothesis beyond QCD (as opposed to the Ellis-Jaffe sum rule⁵ for singlet quantities).
- (2) The estimations of the unmeasured low- x part of the integral are more reliable.
- (3) The pQCD evolution equations are simpler.
- (4) At low Q^2 , the χ PT calculations are more reliable due to the cancellation of the Δ_{1232} contribution.

These advantages might help in extending the validity domains of pQCD and χ PT. It is conceivable that the hadron-parton gap might be bridged, allowing one to fundamentally describe the nucleon structure at all scales⁶ for the first time. Hence the Bjorken sum is arguably one of the most convenient quantities to measure in the resonance region to understand the hadron-parton transition.

Precise data are available from the Thomas Jefferson National accelerator facility (JLab). Results were published on the proton⁷ and deuterium⁸, from CLAS in Hall B and on ^3He from Hall A^{9,10}. We used these data to extract the Bjorken sum from $Q^2 = 0.16$ to 1.1 GeV^2 . To combine proton and neutron data, the ^3He data were reanalyzed at the same Q^2 as those of Ref.⁷. For consistency, the unmeasured low- x part of the integral was re-evaluated for the three data sets using a consistent prescription¹¹. The part beyond the validity range of Ref.¹¹ was estimated using a Regge form and forcing the total integral as measured by the SLAC E155 experiment¹² at $Q^2=5 \text{ GeV}^2$ and completed by the estimation¹¹ and our Regge form, to verify the Bjorken sum rule at $Q^2=5 \text{ GeV}^2$. The results are shown on Figure 1. The elastic contribution is not included. The negative horizontal bands give the systematic uncertainties on the two data sets. Also plotted are the SLAC E143 results in the resonance region¹³. Two χ PT calculations, from Bernard *et al.*¹⁴ and Ji *et al.*¹⁵, are shown at low Q^2 . At $Q^2 = 0$, the Bjorken sum is constrained by the GDH sum rule (see eq. 2). At higher Q^2 the leading twist calculation up to third order in α_s is shown by the gray band. Its width is due to the uncertainty on α_s . The model of Soffer

and Teryaev¹⁶ overestimates the data at large Q^2 . An improved version accounting for the pQCD radiations was presented¹⁷ and seems to agree better with the data. The calculation from Burkert and Ioffe¹⁸ agrees well with the data.

In the particular case of the χ PT calculation done in the heavy Baryon approximation¹⁵, the comparison with the data may indicate that the χ PT domain of validity is indeed extended since they agree, up to about $Q^2 = 0.25 \text{ GeV}^2$ (to be compared to $Q^2 = 0.1 \text{ GeV}^2$ typically for singlet quantities). However, the calculations from Bernard *et al.*¹⁴, that do not employ the heavy Baryon approximation, do not support this conclusion. In any case, a gap between the χ PT calculations and the pQCD calculation clearly remains. More details can be found in Ref. ¹⁹

2. Twist Analysis

It is remarkable that the data agree with the leading twist calculation down to quite low Q^2 . This may indicate that higher twist terms are small or cancel each other. To quantitatively address this, we performed a twist analysis. The coefficient μ_4^{p-n} of the $1/Q^2$ correction to Eq. 1 is:

$$\mu_4^{p-n} = \frac{M^2}{9} (a_2^{p-n} + 4d_2^{p-n} + 4f_2^{p-n}), \quad (3)$$

where a_2^{p-n} is the target mass correction given by the second moment of g_1^{p-n} , and d_2^{p-n} is a twist-3 matrix element given by

$$d_2^{p-n} = \int_0^1 dx x^2 (2g_1^{p-n} + 3g_2^{p-n}). \quad (4)$$

The term a_2^{p-n} is computed using our data and d_2^{p-n} is obtained from SLAC and JLab data²⁰. The twist 4 term f_2^{p-n} is extracted from a fit of the Bjorken sum including the elastic contribution. After re-estimating the low- x part of the world data using our same method, we fit our data together with the world data in the $0.66\text{--}15 \text{ GeV}^2$ Q^2 -range. To verify that the twist series is convergent, the next twist term μ_6^{p-n} was included. We obtain, for $Q^2 = 1 \text{ GeV}^2$, $f_2^{p-n} = -0.17 \pm 0.05(\text{uncor})_{-0.05}^{+0.04}(\text{cor})$ and $\mu_6^{p-n} = 0.09 \pm 0.02(\text{uncor}) \pm 0.01(\text{cor}) \text{ GeV}^4$ where *uncor* (*cor*) specifies the uncertainties due to the point to point uncorrelated (*correlated*) uncertainty on the JLab data. Comparing $\mu_6^{p-n}/Q^4 = 0.09 \pm 0.02$ to $\mu_4^{p-n}/Q^2 \simeq -0.06 \pm 0.02$, we find that the Q^{-2} and Q^{-4} terms have opposite sign and similar magnitude, making the overall twist correction small at $Q^2 = 1 \text{ GeV}^2$. This result may




Figure 1. Q^2 -evolution of the Bjorken sum. The Bjorken sum formed using neutron data extracted from ^3He (D) data is shown by the triangles (squares) and the horizontal band spanning from $Q^2 = 0.16$ to 1.0 GeV^2 (from $Q^2 = 0.35$ to 1.1 GeV^2) is the corresponding systematic uncertainty. The resonance data from SLAC E143 are also shown (circles). The pQCD calculation at leading twist is represented by the gray band. Two χPT calculations (Ji *et al.* and Bernard *et al.*) can be seen at low Q^2 as well as the GDH slope which constrains the Bjorken sum near the photon point. The dashed curves are the predictions of two phenomenological models (Burkert and Ioffe, bottom curve, and Soffer and Teryaev, top curve).

explain why many experimental JLab results tend to indicate that pQCD works down to surprisingly low Q^2 .

3. Summary and outlook

We have extracted the Bjorken sum in the Q^2 range of 0.16 - 1.1 GeV^2 . Compared to singlet quantities, the χPT calculation seems to agree with the data over a larger Q^2 range only in the case of the Heavy Baryon approximation. Such possible improvement is not seen with the calculations of Ref.¹⁴. This last point is not unlike the conclusion reached in Ref.²¹ where

χ PT calculations were compared to measurements of the generalized spin polarizability δ_{LT} , a quantity in which the Δ degrees of freedom are also suppressed. At any rate the parton to hadron gap, if smaller, is not bridged yet. The magnitudes of the higher twists were extracted, in particular f_2^{p-n} . The higher twist effects appear to be small, due to a cancellation of the $1/Q^2$ and $1/Q^4$ terms. The analysis of new proton and deuterium data from CLAS will be finalized soon. These cover a larger Q^2 range (0.05 to 4 GeV²) with improved statistics²². Data at even lower Q^2 are expected to be available in the upcoming years from both Jefferson Lab Hall A (³He and neutron²³) and Hall B (proton)²⁴.

References

1. See for ex. S. Choi, these proceedings.
2. J. D. Bjorken, Phys. Rev. **148** 1467 (1966).
3. M. Anselmino, B.L. Ioffe and E. Leader, Sov. J. Nucl. **49**, 136, (1989).
4. X. Ji and J. Osborne, J.Phys. G27 127 (2001).
5. J. Ellis and R.L. Jaffe, Phys. Rev. **D9** 1444 (1974).
6. V. D. Burkert, Phys. Rev. D **63**, 097904 (2001).
7. R. Fatemi *et al.*, Phys. Rev. Lett. **91**, 222002 (2003).
8. J. Yun *et al.*, Phys. Rev. C **67**, 055204 (2003).
9. M. Amarian *et al.*, Phys. Rev. Lett. **89**, 242301 (2002).
10. M. Amarian *et al.*, Phys. Rev. Lett. **92**, 022301 (2004).
11. N. Bianchi and E. Thomas, Phys. Lett. **B450**, 439 (1999); Nucl. Phys. Proc. Suppl. **82**, 256 (2000).
12. P.L. Anthony *et al.*, Phys. Lett. B **493**, 19 (2000).
13. K. Abe *et al.*, Phys. Rev. Lett. **78**, 815 (1997).
14. V. Bernard, T. R. Hemmert and Ulf-G. Meißner, Phys. Rev. D **67**, 076008 (2003).
15. X. Ji, C. W. Kao and J. Osborne, Phys. Lett. **B472**, 1 (2000).
16. J. Soffer and O. V. Teryaev, Phys. Lett. **B545**, 323 (2002).
17. J. Soffer, these proceedings
18. V. D. Burkert and B. L. Ioffe, Phys. Lett. **B296**, 223 (1992); J. Exp. Theor. Phys. **78**, 619 (1994).
19. A. Deur *et al.* hep-ex/0407007.
20. X. Zheng *et al.*, nucl-ex/0405006, accepted for publication in Phys. Rev. C.
21. M. Amarian *et al.*, nucl-exp/0406005, accepted for publication in Phys. Rev. Lett.
22. G. Dodge, these proceedings.
23. J.P. Chen, A. Deur and F. Garibaldi, JLab experiment E97-110.
24. M. Battaglieri, A. Deur, R. De Vita and M. Ripani, JLab experiment E03-006.